


## IMAGE

# Co-activation of VEGF and NMDA receptors promotes synaptic targeting of AMPA receptors

P De Rossi<sup>1,2,3</sup>, E Harde<sup>4,5,6</sup>, JP Dupuis<sup>7,8</sup>, L Martin<sup>1,2,3</sup>, N Chounlamountri<sup>1,2,3</sup>, M Bardin<sup>1,2,3</sup>, C Watrin<sup>1,2,3</sup>, C Benetollo<sup>1,2,9</sup>, K Pernet-Gallay<sup>10,11</sup>, HJ Luhmann<sup>12</sup>, J Honnorat<sup>1,2,13</sup>, G Malleret<sup>1,2,14</sup>, L Groc<sup>7,8</sup>, A Acker-Palmer<sup>4,5,6</sup>, PA Salin<sup>1,2,14</sup> and C Meissirel<sup>1,2,3</sup>



<sup>1</sup>Institut National de la Santé et de la Recherche Médicale, Unité 1028, Centre National de la Recherche Scientifique, UMR 5292, Lyon, France; <sup>2</sup>Claude Bernard University Lyon 1, Lyon, France; <sup>3</sup>Neurooncology and Neuroinflammation, Lyon Neuroscience Research Center, Lyon, France; <sup>4</sup>Institute of Cell Biology and Neuroscience and BMLS, Goethe University Frankfurt, Frankfurt, Germany; <sup>5</sup>Max Planck Institute for Brain Research, Frankfurt, Germany; <sup>6</sup>Focus Program Translational Neurosciences, University of Mainz, Mainz, Germany; <sup>7</sup>Interdisciplinary Institute for Neuroscience, UMR 5297, Université de Bordeaux, Bordeaux, France; <sup>8</sup>Interdisciplinary Institute for Neuroscience, UMR 5297, Centre National de la Recherche Scientifique, Bordeaux, France; <sup>9</sup>Functional Neurogenomics and Optogenetics, Lyon Neuroscience Research Center, Lyon, France; <sup>10</sup>Grenoble Institute of Neurosciences, Grenoble, France; <sup>11</sup>INSERM U836, Microscopy and Electron Microscopy Platform, Grenoble, France; <sup>12</sup>Institute of Physiology, University Medical Center, University of Mainz, Mainz, Germany; <sup>13</sup>Neuro-Oncology Department, Hospices Civils de Lyon, Hôpital Neurologique, Lyon, France and <sup>14</sup>Forgetting and Cortical Dynamics, Lyon Neuroscience Research Center, Lyon, France